arXiv:math/0502113v1 [math.NT] 6 Feb 2005

q-ANALOGUES OF THE SUMS OF POWERS OF CONSECUTIVE INTEGERS

Taekyun Kim

Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim@kongju.ac.kr (or tkim64@hanmail.net)

Abstract. Let n, k be the positive integers $(k > 1)$, and let $S_{n,q}(k)$ be the sums of the n-th powers of positive q-integers up to $k-1$: $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l l^n$. Following an idea due to J. Bernoulli, we explore a formula for $S_{n,q}(k)$.

1. Introduction

In the early of 17th century Faulhaber [3, 6, 11] computed the sums of powers $1^m + 2^m + \cdots + n^m$ up to $m = 17$ and realized that for odd m, it is not just a polynomial in n but a polynomial in the triangular number $N = n(n+1)/2$. A good account of Faulhaber's work was given by Knuth [3, 6, 11]. In 1713, J. Bernoulli first discovered the method which one can produce those formulae for the sum $\sum_{l=1}^{n} l^k$ for any natural numbers k, cf. [1, 2, 4, 5, 13]. Let q be an indeterminate which can be considered in complex number field, and for any integer k define the q-integer as $[k]_q = \frac{q^k - 1}{q-1}$, cf. [2, 7, 8, 9, 10]. Note that $\lim_{q \to 1} [k]_q = k$. Recently, many authors are studying the problems of q-analogues of sums of powers of consecutive integers [3, 4, 5, 6, 7, 8, 11, 12, 13, 14]. In this paper we consider the q-analogues of the sums of powers of consecutive integers due to J. Bernoulli . For any positive integers $n, k(> 1)$,

---

2000 Mathematics Subject Classification 11S80, 11B68

Key words and phrases: Sums of powers, Bernoulli number, Bernoulli polynomials

Typeset by $\mathcal{AMS}$-TEX

let $S_{n,q}(k)=\sum_{l=0}^{k-1} q^l l^n$. Following an idea due to J. Bernoulli, we explore a formula for $S_{n,q}(k)$ as follows:

$$q^{-k}S_{l-1,q}(k)+(q^{-k}\log q)S_{l,q}(k)=\frac{1}{l}\sum_{i=0}^{l-1}\binom{l}{i}B_{i,q}k^{l-i}+\frac{(1-q^{-k})B_{l,q}(0)}{l},$$

where $B_{i,q}$ are the q-analogue of Bernoulli numbers.

2. q-ANALOGUE OF THE SUMS OF THE n-TH POWERS OF POSITIVE INTEGERS UP TO $k-1$

Let j be the positive integers. Then we easily see that

$$q^{j+1}(j+1)-q^j j=(q-1)q^j j+q^{j+1}. \tag{1}$$

From Eq.(1), we note that

$$(q-1)\sum_{j=0}^{k-1}q^j j+q[k]_q=q^k k. \tag{2}$$

Thus, we have the following:

$$S_{1,q}(k)=\sum_{j=0}^{k-1}q^j j=\frac{q^k k-q[k]_q}{q-1}. \tag{3}$$

Note that $\frac{k(k-1)}{2}=\lim_{q\to 1}\frac{q^k k-q[k]_q}{q-1}=S_1(k)$. By the same method of Eq.(1), we easily see that

$$q^{j+1}(j+1)^2-q^j j^2=(q-1)q^j j^2+2q^{j+1}j+q^{j+1}. \tag{4}$$

Thus, we obtain

$$S_{2,q}(k)=\sum_{j=0}^{k-1}q^j j^2=\frac{q^k k^2}{q-1}-2q\frac{q^k k-q[k]_q}{(q-1)^2}-\frac{q[k]_q}{q-1}. \tag{5}$$

From the simple calculation, we note that

$$\begin{aligned}&q^{j+1}(j+1)^3-q^j j^3=(q-1)q^j j^3+3qq^j j^2+3qq^j j+q^{j+1},\\&q^k k^3=(q-1)S_{3,q}(k)+3qS_{2,q}(k)+3qS_{1,q}(k)+q[k]_q.\end{aligned} \tag{6}$$

Thus, we see that

$$S_{3,q}(k) = \frac{q^k k^3}{q-1} - \frac{3q}{q-1} S_{2,q}(k) - \frac{3q}{q-1} S_{1,q}(k) - \frac{q[k]_q}{q-1}. \tag{7}$$

Let n, k be the positive integers ($k > 1$) and let $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l l^n$. Then we have

$$q^{l+1}(l+1)^n - l^{n+1}q^l = q^{l+1} \sum_{i=0}^{n} \binom{n+1}{i} l^i + (q-1) l^{n+1} q^l.$$

Summing over l from 0 to $k-1$, the left-hand side becomes $q^k k^{n+1}$, but the right-hand side is a linear combination of $S_{i,q}(k)$ as follows:

$$q^k k^{n+1} = q(n+1) S_{n,q}(k) + q \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q}(k) + (q-1) S_{n+1,q}(k). \tag{8}$$

Thus, we obtain the following :

Theorem A. *Let n, $k(>1)$ be the positive integers. Then we have*

$$S_{n,q}(k) = \frac{k^{n+1}}{n+1} q^{k-1} - \frac{1}{n+1} \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q}(k) + \frac{q-1}{q(n+1)} S_{n+1,q}(k). \tag{9}$$

Remark. Note that

$$\begin{aligned} \lim_{q \to 1} S_{n,q}(k) &= \lim_{q \to 1} \left(\frac{k^{n+1}}{n+1} q^{k-1} - \frac{1}{n+1} \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q}(k) + \frac{q-1}{q(n+1)} S_{n+1,q}(k) \right) \\ &= \frac{k^{n+1}}{n+1} - \frac{1}{n+1} \sum_{i=0}^{n-1} \binom{n+1}{i} S_i(k) = S_n(k), \text{ cf. [13].} \end{aligned}$$

3. A formula for $S_{n,q}(k)$

In this section, we assume $q \in \mathbb{C}$ with $|q| < 1$. Now, we consider the q-Bernoulli polynomials as follows:

$$\frac{\log q + t}{qe^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,q}(x) \frac{t^n}{n!}, \text{ see [7, 8].} \tag{10}$$

In the case $x=0$, $B_{n,q}(=B_{n,q}(0))$ will be called the q-Bernoulli numbers, see [8, 9]. Let $F_q(t,x)=\frac{\log q+t}{qe^t-1}e^{xt}$. Then we see that

$$F_q(t,x)=-(t+\log q)\sum_{n=0}^{\infty}e^{(n+x)t}q^n,\ t\in\mathbb{C}\text{ with }|t|<2\pi,\text{ cf.[8, 9, 10].}\tag{11}$$

In [8], it was known that

$$B_{n,q}(x)=\sum_{j=0}^{n}\binom{n}{j}B_{j,q}x^{n-j}=m^{n-1}\sum_{i=0}^{m-1}q^iB_{n,q^m}(\frac{x+i}{m}),\ \text{for } n\geq 0.\tag{12}$$

By (10) and (11), we easily see that the q-Bernoulli numbers can be rewritten as

$$B_{0,q}=\frac{q-1}{\log q},\ q(B_q+1)^k-B_{k,q}=\delta_{k,1},\ \text{for } k\geq 1,$$

where $\delta_{k,1}$ is Kronecker symbol and we use the usual convention about replacing B_q^i by $B_{i,q}$, cf. [8, 9, 10]. From the simple calculation, we note that

$$-\sum_{n=0}^{\infty}e^{(n+k)t}q^n+\sum_{n=0}^{\infty}e^{nt}q^{n-k}=\sum_{l=0}^{\infty}(q^{-k}\sum_{n=0}^{k-1}n^lq^n)\frac{t^l}{l!}=\sum_{l=1}^{\infty}(q^{-k}l\sum_{n=0}^{k-1}n^{l-1}q^n)\frac{t^{l-1}}{l!}.\tag{13}$$

Thus, we have

$$\begin{aligned}\sum_{l=0}^{\infty}(B_{l,q}(k)-q^{-k}B_{k,q}(0))\frac{t^l}{l!}&=-(t+\log q)\sum_{n=0}^{\infty}e^{(n+k)t}q^n+(t+\log q)\sum_{n=0}^{\infty}e^{nt}q^{n-k}\\&=\sum_{l=0}^{\infty}\left(q^{-k}lS_{l-1,q}(k)+(q^{-k}\log q)S_{l.q}(k)\right)\frac{t^l}{l!}.\end{aligned}\tag{14}$$

By comparing the coefficient on both side in Eq.(14), we obtain the following:

Theorem B. *Let l,k be positive integers ($k>1$). Then we see that*

$$q^{-k}S_{l-1,q}(k)+(q^{-k}\log q)S_{l,q}(k)=\frac{B_{l,q}(k)-q^{-k}B_{l,q}(0)}{l}.$$

Remark. Note that

$$S_{l-1}(k)=\lim_{q\to 1}\left(q^{-k}S_{l-1,q}(k)+(q^{-k}\log q)S_{l,q}(k)\right)=\frac{B_l(k)-B_l(0)}{l},\ \text{cf. [13]},$$

where $B_l(k)$ are called ordinary Bernoulli polynomials.

REFERENCES

1. T. Apstol, *Introduction to analytic number theory*, Undergraduate Texts in Math., Springer-Verlag, New York, 1986.
2. L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
3. J. Faulhaber, *Academia Algebrae, Darinnen die miraculosische inventiones zu den höchsten Cossen weiters continuirt und profitiert werden*, Augspurg, bey Johann Ulrich Schönigs, 1631.
4. K. C. Garrett and K. Hummel, *A combinatorial proof of the sum of q-cubes*, Electronic J. Combinatorics **11** (2004), R 9.
5. G. Gasper, M. Rahman, *Basic hypergeometric series*, Encyclopedia of Mathematics and Its Applications, Vol 96, Secod Edition, Cambridge Univ. Press, 2004.
6. V. J. W. Guo, J. Zeng, *A q-analogue of Faulhaber's formula for sums of powers*, arXiv:math.CO/0501441 (2005).
7. T. Kim, *Sums of powers of consecutive q-integers*, Advan. Stud. Contemp. Math. **9** (2004), 15-18.
8. T. Kim, *A new approach to q-zeta function*, arXiv:math.NT/0502005 (2005).
9. T. Kim, *q-Volkenborn Integration*, Russian J. Math. Phys. **9** (2002), 288-299.
10. T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russian J. Math. Phys. **11** (2004), 71-76.
11. D. E. Knuth, *Johann Faulhaber and sums of powers*, Math. Comput. **61** (1993), 277-294.
12. M. Schlosser, *q-analogues of the sums of consecutive integers, squares,cubes, quarts and quints*, Electronic J. Combinatorics **11** (2004), R 71.
13. Y.-Y. Shen, *A note on the sums of powers of consecutive integers*, Tunghai Science **5** (2003), 101-106.
14. S. O. Warnaar, *On the q-analogues of the sums of cubes*, Electronic J. Combinatorics **11** (2004), N 13.